


Bowel Cancer (C18-C20): 2010-2012

Distribution of Cases Diagnosed By Anatomical Site, UK

	Males		Females	
	Average Cases	%	Average Cases	%
Caecum (C18.0)	2,829	12.2%	3,145	17.2%
Appendix (C18.1)	254	1.1%	356	1.9%
Ascending Colon (C18.2)	1,691	7.3%	1,790	9.8%
Hepatic Flexure (C18.3)	673	2.9%	616	3.4%
Transverse Colon (C18.4)	1,098	4.7%	1,148	6.3%
Splenic Flexure (C18.5)	539	2.3%	393	2.1%
Descending Colon (C18.6)	707	3.0%	558	3.0%
Sigmoid Colon (C18.7)	5,380	23.1%	3,740	20.4%
Colon, Overlapping and Unspecified (C18.8-C18.9)	1,153	5.0%	1,243	6.8%
Rectosigmoid Junction (C19)	1,631	7.0%	1,089	5.9%
Rectum (C20)	7,327	31.5%	4,240	23.1%
Total	23,282	100.0%	18,317	100.0%



Source: cruk.org/cancerstats

You are welcome to reuse this Cancer Research UK statistics content for your own work. Credit us as authors by referencing Cancer Research UK as the primary source. Suggested style: Cancer Research UK, full URL of the page, Accessed [month] [year].